


Titre: Title:	Textural analysis of late gadolinium enhanced magnetic resonance images can discriminate acute from chronic myocardial infarction
Auteurs: Authors:	Pascale Beliveau, Farida Cheriet, Stasia A. Anderson, Joni Taylor, Andrew E. Arai, & Li-Yueh Hsu
Date:	2014
Type:	Communication de conférence / Conference or Workshop Item
Référence: Citation:	Beliveau, P., Cheriet, F., Anderson, S. A., Taylor, J., Arai, A. E., & Hsu, L.-Y. (janvier 2014). Textural analysis of late gadolinium enhanced magnetic resonance images can discriminate acute from chronic myocardial infarction [Affiche]. 17th annual SCMR Scientific Sessions, New Orleans, LA, USA (2 pages). Publié dans Journal of Cardiovascular Magnetic Resonance, 16(S1). https://doi.org/10.1186/1532-429x-16-s1-p182

 **Document en libre accès dans PolyPublie**
Open Access document in PolyPublie

URL de PolyPublie: PolyPublie URL:	https://publications.polymtl.ca/10603/
Version:	Version officielle de l'éditeur / Published version Révisé par les pairs / Refereed
Conditions d'utilisation: Terms of Use:	CC BY

 **Document publié chez l'éditeur officiel**
Document issued by the official publisher

Nom de la conférence: Conference Name:	17th annual SCMR Scientific Sessions
Date et lieu: Date and Location:	2014-01-16 - 2014-01-19, New Orleans, LA, USA
Maison d'édition: Publisher:	Springer
URL officiel: Official URL:	https://doi.org/10.1186/1532-429x-16-s1-p182
Mention légale: Legal notice:	© 2014 Beliveau et al.; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

POSTER PRESENTATION

Open Access

Textural analysis of late gadolinium enhanced magnetic resonance images can discriminate acute from chronic myocardial infarction

Pascale Beliveau^{1,2*}, Farida Cheriet², Stasia A Anderson¹, Joni Taylor¹, Andrew E Arai¹, Li-Yueh Hsu¹

From 17th Annual SCMR Scientific Sessions
New Orleans, LA, USA. 16-19 January 2014

Background

Late gadolinium enhanced magnetic resonance (LGE MR) imaging is the current standard modality for evaluating the extent of myocardial damage in the presence of ischemic cardiomyopathies (ICM). However it is difficult to differentiate acute from chronic myocardial infarctions (MI) by standard signal intensity thresholding methods. We hypothesize that textural analysis of myocardial infarct regions has the potential to discriminate acute from chronic MI in high resolution LGE CMR images.


Methods

The experiment was performed on 10 rats (8 month-old males). Five acute MI were created with 30-minute LAD occlusion and imaging was performed 3 hours post-reperfusion. Permanent ligation of the LAD was performed for the 5 chronic MI rats, imaged 2 months after the intervention. LGE MR images were acquired at 50-55 μm isotropic voxel resolution using a 3D gradient echo sequence on a 7T Bruker scanner. Histological staining on short-axis slices of the excised hearts (myoglobin for acute and Masson Trichrome for chronic MI) was the reference for manual segmentation of infarct regions of interest. Three texture features (homogeneity, energy and entropy) were derived from Haralick's grey level co-occurrence matrix and computed in 3D. Student's T-test was performed to compare the two groups.

Results

All texture features were significantly different between acute versus chronic MI ($p < 0.05$). The homogeneity

and energy texture values were 1.40 and 1.43 times higher in chronic than in acute MI. A higher homogeneity or energy value indicates a flatter pixel-to-pixel variation. However, the entropy texture was 1.12 times higher in acute MI which indicates a textural complexity in the infarct regions by the presence of microvascular obstructions. Figure 1 shows example images of acute and chronic MI for the ex vivo rat hearts.



¹National Heart Lung and Blood Institute, National Institutes of Health, Rockville, Maryland, USA
Full list of author information is available at the end of the article

Conclusions

Textural features were significantly different in animal models of acute vs. chronic MI with high resolution LGE CMR imaging. Further study is needed to establish the performance of discriminating MI age with texture features in a clinical setting. Textural analysis, a reproducible method, of myocardial infarct regions in LGE CMR images could potentially improve the quantification and diagnosis of MI in ischemic cardiomyopathies.

Funding

This research was supported by the Intramural Research Program of the National Heart, Lung, and Blood Institute, National Institutes of Health.

Authors' details

¹National Heart Lung and Blood Institute, National Institutes of Health, Rockville, Maryland, USA. ²Ecole Polytechnique of Montreal, Montreal, Quebec, Canada.

Published: 16 January 2014

doi:10.1186/1532-429X-16-S1-P182

Cite this article as: Beliveau *et al.*: Textural analysis of late gadolinium enhanced magnetic resonance images can discriminate acute from chronic myocardial infarction. *Journal of Cardiovascular Magnetic Resonance* 2014 **16**(Suppl 1):P182.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

