

Titre: Title:	P0006: Hearts on Fire
Auteurs: Authors:	Giuseppe Di Labbio, Jérôme Vétel, & Lyes Kadem
Date:	2018
Type:	Communication de conférence / Conference or Workshop Item
Référence: Citation:	Di Labbio, G., Vétel, J., & Kadem, L. (novembre 2018). P0006: Hearts on Fire [Affiche]. 71th Annual Meeting of the APS Division of Fluid Dynamics - Gallery of Fluid Motion, Atlanta, Georgia (1 page). https://doi.org/10.1103/aps.dfd.2018.gfm.p0006

 **Document en libre accès dans PolyPublie**
Open Access document in PolyPublie

URL de PolyPublie: PolyPublie URL:	https://publications.polymtl.ca/10624/
Version:	Version officielle de l'éditeur / Published version Révisé par les pairs / Refereed
Conditions d'utilisation: Terms of Use:	CC BY-NC

 **Document publié chez l'éditeur officiel**
Document issued by the official publisher

Nom de la conférence: Conference Name:	71th Annual Meeting of the APS Division of Fluid Dynamics - Gallery of Fluid Motion
Date et lieu: Date and Location:	2018-11-18 - 2018-11-20, Atlanta, Georgia
Maison d'édition: Publisher:	American Physical Society
URL officiel: Official URL:	https://doi.org/10.1103/aps.dfd.2018.gfm.p0006
Mention légale: Legal notice:	This work is licensed under the Creative Commons Attribution-NonCommercial 4.0 International License. Any reuse must credit the author(s) and provide a link back to this page.

Hearts on Fire

G. Di Labbio^{1,2}, J. Vétel², L. Kadem¹

¹Laboratory of Cardiovascular Fluid Dynamics, Concordia University Montréal, Canada

²Laboratoire de Dynamique des Fluides, Polytechnique Montréal, Canada

The left ventricle is the heart's powerhouse, responsible for pumping oxygen- and nutrient-rich blood to each and every tissue throughout the body. By consequence, diseases affecting this laborious chamber are not only more common but often more serious. The fluid dynamics in the healthy left ventricle is characterized by the impulsive formation of a vortex ring during its filling phase which facilitates the following ejection with very little energy loss. In the case of a leaking aortic valve however, the left ventricle will fill from two sides, resulting in the interaction between two pulsatile jets in a confined elastic vessel, marking a new and unique fluid dynamics problem. Here, we show the backward finite-time Lyapunov exponent in the healthy and diseased left ventricles, revealing the attracting Lagrangian coherent structures associated with the aortic (A) and mitral (M) inflows. The background shows the time-frequency spectra of velocity signals over one cardiac cycle taken at the entrance of the mitral (right) and aortic (left) inflows for the healthy (top) and diseased (bottom) cases. The overlay and perceptually-uniform colormap (thanks to Stéfan van der Walt and Nathaniel Smith) are such that the high frequency bursts (up to 200 Hz) occurring early in the filling phase give the appearance of fire.

