



Titre: Central and peripheral shoulder fatigue pre-screening using the
Title: sigma-lognormal model: a proof of concept

Auteurs: Anaïs Laurent, Réjean Plamondon, & Mickael Begon
Authors:

Date: 2020

Type: Article de revue / Article

Référence: Laurent, A., Plamondon, R., & Begon, M. (2020). Central and peripheral shoulder
fatigue pre-screening using the sigma-lognormal model: a proof of concept.
Citation: Frontiers in Human Neuroscience, 14, 171 (16 pages).
<https://doi.org/10.3389/fnhum.2020.00171>

 **Document en libre accès dans PolyPublie**
Open Access document in PolyPublie

URL de PolyPublie: <https://publications.polymtl.ca/45957/>
PolyPublie URL:

Version: Matériel supplémentaire / Supplementary material
Révisé par les pairs / Refereed

Conditions d'utilisation: CC BY
Terms of Use:

 **Document publié chez l'éditeur officiel**
Document issued by the official publisher

Titre de la revue: Frontiers in Human Neuroscience (vol. 14)
Journal Title:

Maison d'édition: Frontiers Media S.A.
Publisher:

URL officiel: <https://doi.org/10.3389/fnhum.2020.00171>
Official URL:

Mention légale: Copyright © 2020 Laurent, Plamondon and Begon. This is an open-access article
Legal notice: distributed under the terms of the Creative Commons Attribution License (CC BY). The
use, distribution or reproduction in other forums is permitted, provided the original
author(s) and the copyright owner(s) are credited and that the original publication in
this journal is cited, in accordance with accepted academic practice. No use, distribution
or reproduction is permitted which does not comply with these terms.

Supplementary Material

Table 1 | Resume of the parameters extracted and their signification

Input level: central system	
t_0	It is the time that takes the brain to perceive the stimulus and emit the command to the musculoskeletal system. It refers to the moment when a population of neurons sends a motor command, it occurs after the audible stimulus is perceived and the motor command prepared.
$\Delta(t_0)$	It reflects the rhythmicity of an input command. It represents the time elapsed between two successive t_0 and is used in the oscillations only.
D	It corresponds to the distance covered by the resulting lognormal.
θ_s	It is the starting angle of the lognormal.
θ_e	It is the ending angle of the lognormal.
Timing properties of the neuromuscular system: peripheral system	
μ	Also known as the logtime delay, it represents the time taken to reach half of the distance movement on a logarithmic scale. It corresponds to the rapidity of a reaction to a command by a system.
σ	Also known as the logresponse time, it represents the time taken from the neuromuscular system to respond to a command on a logarithmic scale. It is also linked to the movement duration and is a measure of the asymmetry of the lognormal.
Global state of the neuromotor system	
Nblog	It is the number of lognormals required to reconstruct the velocity profile of the movement.
SNR	It is the measure of the quality of the movement reconstruction.
SNR/Nblog	It is a performance criterion and represents the motor control fluency of a gesture. The lognormality principle predicts that the ideal movement converges toward a lognormal profile. When the SNR/Nblog is higher, the movement is more similar to the ideal one, as postulated by the lognormal behavior.

Motor program execution ($\Delta(t_0)$ is used instead of t_0 for the calculus in the oscillations)

Mode

It is the time at which the maximum value of the lognormal impulse response is reached.

$$M = t_0 + e^{\mu - \sigma^2}$$

Median

It corresponds to the time at which the half value of the integral under the lognormal curve (50% of the covered distance) is reached.

$$m = t_0 + e^{\mu}$$

Time delay

It represents the rapidity of a neuromuscular system to respond to a command.

$$\bar{t} = t_0 + e^{\mu + 0.5\sigma^2}$$

Response time

It is a measure of the spread of the impulse response.

$$s = (\bar{t} - t_0) \sqrt{(e^{\sigma^2} - 1)}$$

Asymmetry

It characterizes the shape of the lognormal.

$$A_c = 1 - e^{-\sigma^2}$$

Other parameters

Reaction time

It is the time needed to start the movement after a stimulus. In the present study, it was computed as the time required to reach 10% of the maximal velocity during the test

Command propagation

It is the duration of the command propagation

$$CP = RT - t_0$$
