

Titre: Risk assessment of drinking water intake contamination from agricultural activities using a Bayesian network
Title:

Auteurs: Raja Kammoun, Natasha McQuaid, Vincent Lessard, Eyerusalem Adhanom Goitom, Michèle Prévost, Françoise Bichai, & Sarah Dorner
Authors:

Date: 2023

Type: Article de revue / Article

Référence: Kammoun, R., McQuaid, N., Lessard, V., Goitom, E. A., Prévost, M., Bichai, F., & Dorner, S. (2023). Risk assessment of drinking water intake contamination from agricultural activities using a Bayesian network. PLOS Water, 2(7), 25 pages.
Citation: <https://doi.org/10.1371/journal.pwat.0000073>

Document en libre accès dans PolyPublie

Open Access document in PolyPublie

URL de PolyPublie: <https://publications.polymtl.ca/54820/>
PolyPublie URL:

Version: Révisé par les pairs / Refereed

Conditions d'utilisation:
Terms of Use:

Document publié chez l'éditeur officiel

Document issued by the official publisher

Titre de la revue: PLOS Water (vol. 2, no. 7)
Journal Title:

Maison d'édition: Public Library of Science
Publisher:

URL officiel: <https://doi.org/10.1371/journal.pwat.0000073>
Official URL:

Mention légale: © 2023 Kammoun et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.
Legal notice:

RESEARCH ARTICLE

Risk assessment of drinking water intake contamination from agricultural activities using a Bayesian network

Raja Kammoun^{1*}, Natasha McQuaid¹, Vincent Lessard², Eyerusalem Adhanom Goitom¹, Michèle Prévost¹, Françoise Bichai¹, Sarah Dorner¹

1 Department of Civil, Geological and Mining Engineering, Polytechnique Montréal, Montreal, Quebec, Canada, **2** Conseil des bassins versants des Mille Îles, Saint-Eustache, Quebec, Canada

* raja.kammoun@polymtl.ca



Abstract

Agricultural activities can result in the contamination of surface runoff with pathogens, pesticides, and nutrients. These pollutants can enter surface water bodies in two ways: by direct discharge into surface waters or by infiltration and recharge into groundwater, followed by release to surface waters. Lack of financial resources makes risk assessment through analysis of drinking water pollutants challenging for drinking water suppliers. Inability to identify agricultural lands with a high-risk level and implement action measures might lead to public health issues. As a result, it is essential to identify hazards and conduct risk assessments even with limited data. This study proposes a risk assessment model for agricultural activities based on available data and integrating various types of knowledge, including expert and literature knowledge, to estimate the levels of hazard and risk that different agricultural activities could pose to the quality of withdrawal waters. To accomplish this, we built a Bayesian network with continuous and discrete inputs capturing raw water quality and land use upstream of drinking water intakes (DWIs). This probabilistic model integrates the DWI vulnerability, threat exposure, and threats from agricultural activities, including animal and crop production inventoried in drainage basins. The probabilistic dependencies between model nodes are established through a novel adaptation of a mixed aggregation method. The mixed aggregation method, a traditional approach used in ecological assessments following a deterministic framework, involves using fixed assumptions and parameters to estimate ecological outcomes in a specific case without considering inherent randomness and uncertainty within the system. After validation, this probabilistic model was used for four water intakes in a heavily urbanized watershed with agricultural activities in the south of Quebec, Canada. The findings imply that this methodology can assist stakeholders direct their efforts and investments on at-risk locations by identifying agricultural areas that can potentially pose a risk to DWIs.

OPEN ACCESS

Citation: Kammoun R, McQuaid N, Lessard V, Goitom EA, Prévost M, Bichai F, et al. (2023) Risk assessment of drinking water intake contamination from agricultural activities using a Bayesian network. *PLOS Water* 2(7): e0000073. <https://doi.org/10.1371/journal.pwat.0000073>

Editor: Soni M. Pradhanang, University of Rhode Island, UNITED STATES

Received: October 31, 2022

Accepted: June 21, 2023

Published: July 27, 2023

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pwat.0000073>

Copyright: © 2023 Kammoun et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data are available at Open Science Framework: https://osf.io/zyekt/?view_only=01527a99a3374a2ab7879e42ff01052b Identifier: DOI [10.17605/OSF.IO/ZYEKT](https://doi.org/10.17605/OSF.IO/ZYEKT).

Funding: This work was supported by the NSERC PURE CREATE program (528220-2019 to RK) and the partner municipalities (3240067-3240072 to SD). The NSERC PURE CREATE had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The partner municipalities had no role in study design, analysis, or preparation of the manuscript. They were only involved in data collection and the decision to publish.

Competing interests: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

1. Introduction

Agricultural practices are significant sources of chemical and microbial pollution to surface waters [1–5], posing a threat to public health. Runoff from agricultural fields and infiltration into groundwater leads to the transport of pesticides and nutrients into receiving waters [6]. The potential for contamination of surface waters depends on the type, quantity, adsorption, and absorption capacity of applied pesticides, nitrogen and phosphorus [7–9]. In addition, soil characteristics [10], rainfall rates [11–13], and agricultural management practices [14–16] are essential factors that determine the susceptibility of runoff to contamination. An increase in nutrient levels from animal and crop production can contribute to the proliferation of cyanobacteria in drinking water sources [17–20]. Some cyanobacteria blooms can release toxins that pose challenges in terms of drinking water safety [21–26]. Animal excrements and manure are also a source of microbial contamination in surface waters [27–30]. Factors influencing the survival and transport of pathogens in water resources include: water temperature, pH, biotic interactions, manure types and characteristics, and farm management practices [31].

Risk assessments can raise awareness of the threats posed by agricultural contamination to drinking water sources. Several methods to assess the risks associated with agricultural activities are available. In recent years, many risk assessment techniques have used deterministic modeling approaches focused on ecological, ecotoxicological, and human health issues [3, 32–37]. Deterministic modeling is costly, requires extensive calibration, and needs a large amount of site-specific data on water quality and hydrodynamics. The central problem with this approach is that it cannot be implemented by all drinking water suppliers due to a lack of financial resources and data. On the other hand, the deterministic approach could lead to an erroneous risk management decision, as it evaluates the impact of only one risk scenario at a time. Some of these limitations can be addressed using a probabilistic approach that simultaneously considers many risk assessment factors, such as various threat exposure scenarios, uncertainty of the different input variables, as well as uncertainty in the relationship between the parameters defining the system and the risk magnitude [38].

Bayesian networks (BN) are a valuable tool for conducting a probabilistic risk evaluation under uncertainty, and integrating different types of knowledge, such as quantitative and qualitative data, expert opinion, and prior experiences [39]. BNs have been extensively applied in different fields, such as in the fields of ecological risk assessment [40–43], human health risk assessment [40–44], climate change risk assessment [45], disaster risk assessment [46, 47], water quality and treatment [48, 49], best management practices of non-point source pollution [50], and water resource management [51]. To the best of our knowledge, these approaches have not been applied to vulnerability assessments of drinking water sources in agricultural watersheds.

The aim of this study was to develop a probabilistic model that decision-makers could use to swiftly identify animal and crop production areas that generate high chemical and microbial risks at DWIs. Our specific objectives were to: (1) develop a BN that combines the three components of risk assessment: hazard, exposure, and vulnerability; (2) test the applicability of existing methods used to define the conditional independence relationship among variables constituting BNs for drinking water risk assessments; (3) develop a novel method to define an efficient Bayes independence of the input variables for water resource risk assessments; (4) assess the probability distribution of the overall hazard of agricultural activities to pinpoint the drainage basins that require more detailed investigation for source water protection.

2. Materials and methods

2.1 Study area

The study was conducted for four DWIs, serving a population of approximately 440,000, drawing water from a river in Southern Quebec, Canada (Fig 1). Drinking water intakes 1 (DWI_1)




Fig 1. Study site. The sources of the mapped data are as follows: (1) drinking water intakes was acquired from municipal partners, (2) land use data was supplied by the Ministère des Affaires municipales et de l'Habitation of Quebec (MAMH), and (3) the base map utilized was created in the "ArcGIS" software program using the "Basemap" function whose shapefile is noted "World Topographic Map".

<https://doi.org/10.1371/journal.pwat.0000073.g001>

and 4 (DWI_4) are located in the most upstream and downstream sections of the river, respectively. DWI_2 and DWI_3 are located in the middle of the river reach. This river drains a lake fed by the Ottawa River, as described in Jalliffier-Verne et al. [52]. The Ottawa River receives runoff from the largest (146,300 km²) watershed in Eastern Canada, and its average discharge is around 1,950 m³/s [53]. This watershed is dominated by forests (75%) of which 40% is dense mixed wood. Agricultural activities take place over 6% of this watershed surface area. The adjacent watersheds that have a local influence on the 42-km long river that supplies the DWIs of this study cover an area of 1,008 km². This area is zoned as follows: 30% agricultural, 28.69% urban, 20.63% forest land.

Delineating protection zones is a critical step to assess non-point source pollution risks to drinking water sources, reduce the risk of contamination, and support drinking water safety. In this study, the delineation of immediate and intermediate protection areas was carried out to specify the land area that contributes water to the drinking water supply. The immediate protection area is a zone close to a DWI where pollution sources are deemed of concern and where contaminants may be readily found in the raw water with minimal dilution [54]. The intermediate protection area is a different zone where the travel time to the DWI is too short to allow intervention in the case of an accidental spill or a spike in contaminant concentration [54].

In Quebec, provincial regulation requires the delineation of surface water protection zones using a fixed distance approach [54]. This approach comprises only a strip of land on both sides of a river and does not include land surfaces whose relief leads to the drainage of land, lakes, and streams upstream of the river under study. For more information on regulatory methods for delineating surface water protection areas, see the S1 Text. To control diffuse

pollution, it is necessary to consider the drained non-point source pollution lands upstream of the tributary outlets included in the protection zones of the DWI. Therefore, this study develops an improved methodology of delineation based on levels 1 and 2 sub-watersheds. Level 1 sub-watersheds are the large watersheds whose outlet is the study site river. Level 2 sub-watersheds are those that drain into level 1 sub-watersheds and have the river in level 1 sub-watersheds as their outlet.

The geographic information system software (ArcGIS) was used to delineate the protection zones using the following data layers: location of DWIs, topological hydrographic network lines, and high-water mark. The immediate protection zone was delimited within 50 m downstream and 500 m upstream of the water intake as required by Quebec regulation [54]. This area includes any surface water, portions of tributaries, and a 10-m strip of land measured from the high-water mark. Surface and water portions of tributaries were also included in the intermediate protection zone and incorporate a 120-m strip of land measured from the high-water mark. This protection zone was delimited within 50 m downstream and 10 km upstream of the DWI [54]. This delineation was used to identify all level 1 and 2 sub-watersheds with outlets within the boundaries of these protection zones and to include them in the study site. Every sub-watershed selected is a drainage basin (DB) (S1 Fig). We define a drainage basin as a territory including a set of parcels used for one or more anthropogenic activities presenting a source of diffuse pollution and whose drained water is discharged into a watercourse through an outlet. Each drainage basin presents a potential source of microbial and/or chemical contamination to the studied DWIs.

2.2 Bayesian network framework

A probabilistic and spatial risk assessment of agricultural activities impacting DWIs was performed at a regional scale using a BN and ArcGIS. The risk assessment framework was built to integrate the three components of risks: hazard, vulnerability, and exposure (Fig 2).

A BN is a hierarchical representation and a probabilistic graphical model representing a relationship between a range of variables. Every single variable is represented as a node, and relationships between variables are presented by arcs defined by conditional dependencies [55]. This directed, acyclic graph allows the modeling of various uncertain events, facts, and systems [56]. More definitions of BN can be found in other studies [38, 47, 50, 57–60]. Bayesian networks are also known as conditional probability networks, casual probability networks, probabilistic networks, belief networks, Bayesian belief networks, or Bayesian reliability networks.

In this study, the BN was developed using the software Netica 6.09 (Norsys Software Corp), applying the Bayes' theorem [61]. This software is an appropriate tool for creating, changing, and saving networks, as reviewed by Uusitalo [39].

2.2.1 Defining the nodes. The first step to structure the BN model was defining a set of random variables (nodes) and their states, which can be discrete or continuous. Discrete variables have a limited set of fixed values or states that they can take. Continuous variables are classified as real numbers since they may have any value within a certain range [62]. Continuous variables were discretized by breaking them up into intervals by assigning thresholds to define each interval. Input nodes are labeled “parent” nodes, and “child” nodes have input connections from one or more parent nodes [40, 58].

The determination of the critical nodes constituting the BN was carried out based on the problem and the objective of the study, which considers the DWI as the main subject of the risk assessment from agricultural activities. These nodes were chosen to represent all components of the risk assessment model based on the interaction of vulnerability to microbiological




Fig 2. Risk assessment process.

<https://doi.org/10.1371/journal.pwat.0000073.g002>

contamination (VMC), vulnerability to chemical contamination (VCC), vulnerability to water scarcity (VWS), exposure (EXP), and probability of hazard (PH). In the Working Group II (WGII) sixth assessment report from Intergovernmental Panel on Climate Change [63], the hazard is described in terms of the potential occurrence of the chemical and microbial contaminants that may cause degradation of water resource. The exposure depends on the presence of DWI in locations that could be adversely affected. The vulnerability is evaluated as the propensity or predisposition of the DWI to be adversely affected.

In this study, the intrinsic vulnerability of the intake was assessed by examining the sensitivity of the raw water to potential pollutants and the physical characteristics of the intake. Three key variables were used to assess the vulnerability to microbiological contamination of water intakes: (i) the geometric mean of *Escherichia coli* concentrations (GMEC) was used to present the central tendency of the microbial contamination indicator while being robust to extreme values; (ii) the 99th percentile of *E. coli* concentrations (PKEC) was used to account for the peaks of microbial contamination that are usually responsible for waterborne diseases [64]; and (iii) the seasonal variability of microbial contamination (SVMC) that was calculated based on a consecutive disparity index that estimates the variability of *E. coli* concentrations while considering the order of their chronological distribution, and without being independent of the data set's average [65]. All variables were calculated for the four seasons of the year [66] (Winter, Spring, Summer, and Autumn) to study microbial contamination distribution and evaluate the sensitivity to seasonal variations.

Assessment of the vulnerability to chemical contamination was carried out based on the total phosphorus in raw water, which is routinely analyzed at drinking water treatment plants in Quebec. As data for other chemicals, such as nitrate and nitrogen, were not available, the vulnerability to chemical contamination was also estimated by analyzing the land use of activities that may release chemical contaminants in the immediate and intermediate DWIs' protection zones. Raw water quality data available over a period of five years (January 2013 to December 2017) was used to assess the microbiological and chemical vulnerability of DWIs as required by Water Withdrawal and Protection Regulation [54].

The results of the vulnerability to water scarcity from the Levesque [67] study were used to assess vulnerability based on (i) the water levels in the studied river, which depend on the physical characteristics of the DWI and (ii) the water demand assigned to each catchment.

The exposure was estimated based on the location of the outlet of each drainage basin in relation to the intake and its protection areas. This node allows assessing the non-point source pollution exposure pathways to the raw water supply.

The probability of hazard was estimated based on the proportion of the drainage basins occupied by crop and animal production activities. Crop classes and animal classes were used as input variables to estimate surface water contamination levels by pesticides and microbial contaminants.

The final node presents the overall risk probability distribution across three possible states (low, medium, and high) of agricultural activities occupying a watershed located upstream of a DWI by combining vulnerability, exposure, and hazard as outlined below.

The input variables, their states, and the discretization methods are described in Table 1. Once the nodes were set up, links (dependencies) between them were defined using conditional probability tables (CPTs) (Section 2.3.2). Finally, the values for each input parameter were computed using the data associated with the study site to produce a probability distribution for each state associated with the nodes.

The determination of states for the remaining nodes and directed links was accomplished based on scientific knowledge (the literature), the Quebec regulations, and the availability of data in provincial and federal databases. The choice of nodes was discussed with experts and stakeholders to provide a suitable BN, rectify, refine, and fill the potential gaps associated with the variables and their status.

2.2.2 Filling the CPTs. Filling the CPTs of the Bayesian network is the most challenging step in setting up the framework. To accomplish this step, different known methods were tested: (i) expert judgment [38, 58, 59], (ii) the interpolation method [88], and (iii) the Cain method [89, 90]. Methods that did not generate reliable outcomes in terms of probabilities were rejected under the expert decision.

It was challenging to estimate the quantity of chemical and microbiological contaminants released during agricultural production activities, as well as their diffusion, dispersion, and advection into the DWIs. Thus, employing expert judgment and scientific knowledge from the literature were the best methods for determining the conditional distribution of the quantities of pollutants that may be released in the immediate and intermediate protection zones. The exposure to pollutants released by upstream agricultural activities was also assessed using this method. In these situations, completing the CPTs was based on experts' judgments and their interpretation by the extensive research team (rather than through formal interviews). This is due to the nature of the project: experts' judgments were interpreted throughout four years of work on a collaborative project between six municipal partners, a watershed organization, and a research team at Polytechnique Montreal. Thus, the probabilities for each status of each child node were elucidated through discussion meetings with researchers, professionals, and municipal water stakeholders and were enhanced by scientific knowledge.

Table 1. Summary of input variables' properties.

Parent node			Risk component	State discretization methodology	States (from lowest to highest level)	Data source
Name	Description	Type of discretization				
DBA	Drainage basin (DB) area	Continuous	Hazard	Statutes were defined based on percentiles of the area of 4723 sub-watersheds (level 2) in Quebec.	0 to 5 Km ² / 5 to 12 km ² / 12 to 45 km ² / > = 45 km ²	Quebec portal of knowledge on water [68]
PPCO	Plant production cover	Continuous	Hazard	Statutes were defined based on Water Withdrawal and Protection Regulation (WWPR) (schedule IV)	0 to 20% / 20 to 50% / > = 50%	Mapping of land use in the St. Lawrence Lowlands [69]
APCO	Animal production cover	Continuous	Hazard	[54]: land use-based method for assessing water vulnerability.	0 to 20% / 20 to 50% / > = 50%	Data are provided by the <i>Ministère des Affaires municipales et de l'Habitation</i> of Quebec (MAMH).
CPC	Crop production classes	Discrete	Hazard	Environmental impacts depend on the crop class. The statutes were specified based on the data layer available in the ministry database (<i>Ministère de l'environnement et de la lutte contre les changements climatiques (MELCC)</i>), and their classification was made based on the following references: Beaudin et al. [70], Michaud et al. [71], Gasser et al. [72], Giroux [73], and Giroux [74].	No crop production / Non-cultivated agricultural area / Undefined culture / Specialized culture / Perennial culture / Annual crop	Quebec land use [75]
APC	Animal production classes	Discrete	Hazard	Animal manure is a source of microbial contamination for surface waters and contains pathogens. These pathogens are specific to the types of animals from which manure is derived. The statutes were specified based on the data layer available in the MAMH database, and their classification was made based on the following references: Soupir et al. [76], Jaffrezic et al. [77], Lalancette et al. [78], and Haack et al. [79]	No animal production / Class VII ⁽²⁾ / Classes V ⁽³⁾ -VI ⁽⁴⁾ / Classes II ⁽⁵⁾ - III ⁽⁶⁾ -IV ⁽⁷⁾ / Class I ⁽⁸⁾	Data are provided by MAMH.
PZ	Protection zone	Discrete	Exposure	Statutes were defined based on WWPR (Art. 70, 72, and 74) [54]: protection zone delineation.	Extended protection zone / Intermediate protection zone / Immediate protection zone	Data are provided by the watershed organization.
GMEC_X ⁽¹⁾	Geometric mean of <i>E. coli</i> concentrations	Continuous	Vulnerability	Statutes were defined based on the regulation of the drinking water quality [80], and the development of an index of bacteriological and physicochemical quality (IQBP) [81].	0 to 15 CFU/100 ml 15 to 150 CFU/100 ml > = 150 CFU/100 ml	Raw water quality data are provided by drinking water treatment plants managers.
PKEC_X ⁽¹⁾	99th percentile of <i>E. coli</i> concentrations	Continuous	Vulnerability		0 to 200 CFU/100 ml 200 to 2000 CFU/100 ml > = 2000 CFU/100 ml	
SVMC_X ⁽¹⁾	Seasonal variability of microbial contamination	Continuous	Vulnerability		0 to 1 / 1 to 1.5 / > = 1.5	
VWS	Vulnerability to water scarcity	Discrete	Vulnerability	Statutes were defined based on the water scarcity vulnerability index developed by Leveque [67].	Low / Medium / High	Leveque [67] study

(Continued)

Table 1. (Continued)

Parent node			Risk component	State discretization methodology	States (from lowest to highest level)	Data source
Name	Description	Type of discretization				
PKTP	99 th percentile of total phosphorus concentrations	Continuous	Vulnerability	Statutes were defined based on the WWPR [54], and the development of an index of bacteriological and physicochemical quality (IQBP) [81].	0 to 50 µg/l 50 to 100 µg/l / > = 100 µg/l	Raw water quality data are provided by drinking water treatment plants managers.
MTP	Median of total phosphorus concentrations	Continuous	Vulnerability		0 to 30 µg/l 30 to 50 µg/l > = 50 µg/l	
AIMPZ	Area of immediate protection zone	Continuous	Vulnerability		0 to 1km ² / 1 to 5 km ² /	Drinking water intakes vulnerability assessment reports [82–87]
AINPZ	Area of intermediate protection zone	Continuous	Vulnerability	Statutes were defined on the basis of the WWPR [54].	0 to 10 km ² / 10 to 30 km ² / > = 30 km ²	
PIMPA / PINPA	Percentage of the immediate/ intermediate protection zone used for activities that may release chemical contaminants	Discrete	Vulnerability		No / 0 to 20% / 20 to 50% / > = 50%	

⁽¹⁾X refers to season: Winter/ Spring/ Summer/ Autumn

⁽²⁾ Class VII: other types of animal production/ experimental farm/ other agricultural activities; ⁽³⁾ Class VI: apiculture; ⁽⁴⁾ Class V: poultry and egg production; ⁽⁵⁾ Class IV: sheep and goat farming; ⁽⁶⁾ Class III: pig farming; ⁽⁷⁾ Class II: equine farming; ⁽⁸⁾ Class I: beef farming/ dairy cattle/ grazing land and pasture

<https://doi.org/10.1371/journal.pwat.0000073.t001>

However, it is challenging to ensure that the definition of probability distributions of several tables is coherent with each other rather than using simply the judgment of experts. This is mostly because experts lack the rigid regularity of machines [88, 91]. During a long process, boredom and fatigue are enough to introduce errors, uncertainties, and a lack of standardization in the distributions.

One solution would be to standardize the method of obtaining probabilistic information from the experts by providing a much smaller selection of probability distributions. Therefore, the interpolation method was used to obtain the remaining probability distributions in a reasonable time [88]. This method requires assigning a weight to each parent node to quantify the relative strength of its influence over the child node. This step was done based on expert judgment. The sum of the weights must equal one. Then, conditional probability distribution (CPD) elicitation was done for three cases: when all parents nodes are in extreme states (high and low states) and medium states. The determination of the probability distribution for the remaining combinations of the various parents' states was done by interpolation. This method was allied to all CPTs of the Bayesian network. However, it was deemed to be not adequate for our study because an examination of the probability distributions revealed that the resulting probabilities were inconsistent. It was difficult to maintain consistency between the probability distributions, especially in cases where a parent node's state was highly different from the one used for interpolation. In this instance, this method does not account for the influence of variation bias in the interpolation. Therefore, the interpolation method was not pursued further.

The Cain method was also tested. It involves determining CPDs for cases where the relationships between the parent nodes are known; these obtained conditional probability distributions are termed anchor CPDs. Anchors were defined according to the number of parents and the number of statuses associated with each parent. The distribution of missing conditional probabilities was derived by interpolating those corresponding to the anchor CPDs and using interpolation factors. The latter were calculated using the CPDs at anchors. Interpolation

factors of a parent node were calculated for each change from one state to another to quantify the positive to negative state reduction. The processes of defining anchors and calculating interpolation factors are explained in more details in Cain [89] and Mkrtchyan et al. [90].

Only two CPTs, relating to crop and animal activity levels, were completed using the Cain approach because this method is applicable for CPTs in which the effect of the parent state on the child state has a constant direction: the increase in the area occupied by animal and crop production activities results in an increase in the level of agricultural activities in the drainage basin. The results of filling in the probability tables for the other child nodes were judged to be unacceptable since the interpolation is dependent on the number of favourable and unfavourable states for the child state of interest. As a result, there will be the same probability distributions for different child nodes even though the parent nodes have varying degrees of influence on them [92].

Therefore, a novel approach to filling CPTs was developed that combines a mixed aggregation method [93] with expert judgment. The mixed aggregation was used to aggregate indicators in the ecological assessments field using a deterministic approach [93]. It was adapted to be applied in the case of a probabilistic approach and for water resource assessment. First, the different state of each parent node (P_n) was standardized to a common scale from 0 (best possible state) to 1 (worst possible state). Second, the assignment of weights (W_n) for each parent node (P_n) was performed to reflect its importance relative to the child node. Third, causal dependencies were performed with aggregation. In this step, additive and maximal aggregation were combined (Eqs (1), (2), (3), (4), (5)). Additive aggregation (f_{add}) compensates for the poor status of a parent node by other nodes with a good status. On the other hand, maximal aggregation (f_{max}) leads to pessimistic results since only the parent node with the worst status is considered. The combination of the additive and maximal aggregation allows for some compensation while penalizing the very poor state to not neglect an impact that requires more attention to protect water resources.

$$f_{add-max}(P_1, P_2, \dots, P_n) = \alpha f_{add}(P_1, P_2, \dots, P_n) + (1 - \alpha) f_{max}(P_1, P_2, \dots, P_n) \quad (1)$$

Where:

$$f_{add}(P_1, P_2, \dots, P_n) = \sum_{i=1}^n W_i P_i = W_1 P_1 + W_2 P_2 + \dots + W_n P_n \quad (2)$$

$$\sum_{i=1}^n W_i = 1 \quad (3)$$

$$f_{max}(P_1, P_2, \dots, P_n) = \max(P_1, P_2, \dots, P_n) \quad (4)$$

$$\alpha + (1 - \alpha) = 1 \quad (5)$$

α and $(1 - \alpha)$ were chosen according to the relative importance to be given to each aggregation method. The more alpha tends to 0, the more the maximum aggregation is emphasized. Otherwise, additive aggregation tends to be prioritized.

The most favorable weight (W_n) set of parent nodes and aggregation method was determined by performing a calibration that requires knowledge and experience and by building on previous work [82–87] conducted to assess the potential risk of agricultural activities on water resources using a deterministic approach. This step was performed by analyzing the outputs resulting from each possible combination of α and W_n . After validating the most appropriate variables and depending on the number of parent nodes, a 2D or 3D graph was generated to extract the CPDs. These data were used to fill the CPTs. The weights selection and graph

Table 2. Overview of the CPTs filling methods.

Child node			Parent node	Weight of parent node	α	Nominated CPTs filling method	Tested and rejected CPTs filling methods
Name	Description	Risk component					
CAL	Crop activities level	Hazard	PPCO DBA	NA ^(*)	NA	Cain method	<ul style="list-style-type: none"> • Expert judgment • Interpolation method • Mixed aggregation method combined to expert judgment
AAL	Animal activities level	Hazard	APCO DBA				
CRIM	Contaminants that could be released in the immediate protection zone	Vulnerability	PIMPA AIMPZ	NA	NA	Expert judgment	<ul style="list-style-type: none"> • Cain method • Interpolation method • Mixed aggregation method combined to expert judgment
CRIN	Contaminants that could be released in the intermediate protection zone	Vulnerability	PINPA AINPZ				
EXP	Exposure	Exposure	PZ				
CAH	Crops activities hazard	Hazard	CAL CPC	0.9 0.1	1	Mixed aggregation method combined to expert judgment	<ul style="list-style-type: none"> • Interpolation method • Mixed aggregation method
APH	Animal production hazard	Hazard	AAL APC	0.9 0.1	1		
PH	Probability of hazard	Hazard	APH CAH	0.6 0.4	1		
MQRW_ (x) ^(**)	Microbiological quality of raw water	Vulnerability	PKEC_(x) GMEC_(x) SVMC_(x)	0.5 0.4 0.1	0.8		
VMC	Vulnerability to microbiological contamination	Vulnerability	MQRW_W MQRW_SP MQRW_S MQRW_A	0.25 0.25 0.25 0.25	0.5		
RWQLTP	Raw water quality level for total phosphorus	Vulnerability	PKTP MTP	0.2 0.8	0.7		
CVLU	Chemical vulnerability level based on land use	Vulnerability	CRIM CRIN	0.7 0.3	0		
VCC	Vulnerability to chemical contamination	Vulnerability	RWQLTP CVLU	0.8 0.2	0.7		
DWIVL	DWI vulnerability level	Vulnerability	VMC VWS VCC	0.4 0.4 0.2	0.4		
GRL	Global risk level	Risk	PH DWIVL EXP	0.8 0.1 0.1	1		

(*) Not applicable

(**) x refers to season: winter/ Spring/ Summer/ Autumn

<https://doi.org/10.1371/journal.pwat.0000073.t002>

generation process were codified in R (V.1.1.456, RStudio, Inc.). [S1 Table](#) presents an example of CPT associated with the “Global Risk Level (GRL)” node that was determined by combining three parent nodes: probability of hazard (PH), exposure (EXP), and DWI vulnerability level (DWIVL). All probability distributions were calculated based on the 3D graph ([S2 Fig](#)) generated by combining all possible cases related to the parent nodes.

[Table 2](#) shows the methods applied to fill 18 CPTs. This table also lists the methods that were tested out but were rejected because of unreliable outcomes as explained above.

2.2.3 Sensitivity analysis. A sensitivity analysis was performed to quantify the magnitude of the sensitivity of the changes that occur in the endpoint probabilities when the parent nodes are changed and to identify which of the variables are most relevant [40, 60]. This analysis is necessary to understand where more information is needed to reduce uncertainty.




Fig 3. Bayesian network structure for assessing drinking water intake contamination risk from agricultural activities. The parent nodes in gray are associated with input variables that define the properties of a specific drainage basin and variables related to a selected intake.

<https://doi.org/10.1371/journal.pwat.0000073.g003>

This sensitivity analysis was executed by calculating the mutual information (I) between two discrete variables X and Y [94]:

$$I(X, Y) = H(X) - H(X|Y) = \sum_{x \in X} \sum_{y \in Y} p(x, y) \log_2 \frac{p(x, y)}{p(x)p(y)} \quad (6)$$

where the sum is over all states x and y of variables X and Y, respectively, H(X) is the entropy reduction of X before any new finding, H(X|Y) is the conditional entropy of X at given Y. In the case of computing the I between continuous random variables, the double sum has to be replaced by a double integral.

Entropy reduction refers to how much a target node's entropy is decreased when the values of its parent nodes are held constant [95–97]. The amount of this entropy reduction is used to gauge how much the parent nodes, also known as findings nodes, influence the target node. Entropy is determined using the probabilities given to the potential states of the target node. It is a measure of uncertainty or randomness associated with the distribution of probabilities in a system. According to this theory, the target node is thought to be more affected by nodes with higher entropy reduction than nodes with lower entropy reduction [95–97].

These measures of entropy reduction and mutual information between various variables and a given endpoint node were conducted with the “sensitivity to finding” command in the Netica software.

3. Results

3.1 Bayesian network model

The developed BN model integrates 43 nodes (Fig 3) and 1326 conditional probabilities. The nodes constituting this BN are described in Tables 1 and 2. The probabilities associated with




Fig 4. The probability distribution of different categories of vulnerability (global, microbiological contamination, chemical contamination, and scarcity) for all drinking water intakes (DWIs).

<https://doi.org/10.1371/journal.pwat.0000073.g004>

each node whose sum is equal to 100% are shown on the right side of the status window of the detailed BN (S3 Fig). Grey nodes represent the input variables.

3.2 DWIs vulnerability

The findings related to the vulnerability nodes, which are a component of risk, are presented in this section. As illustrated in Fig 4, more than 60% of DWIs were extremely vulnerable. This is most likely because these intakes are in a highly urbanised watershed with agricultural activity (section 2.1). The vulnerability of source water to microbiological contamination was noted at the four intakes. The seasonal variation in microbial contamination of raw water was investigated (S4 Fig), and our results revealed that all intakes were associated with medium to high probability distributions for all seasons. The probabilities describing poor water quality could be due to the transport of microbial contaminants during the late winter and early spring snowmelt period, but also during episodes of heavy rainfall in the summer, and autumn as demonstrated by Sylvestre et al. [29].

DWI_3 and DWI_4 were highly vulnerable to chemical contamination (82.3% and 72.2% probability in the high state, respectively) (Fig 4) as a result of total phosphorus leading to lower quality of the raw water. In addition, a significant area of the intermediate zone of DWI_3 was occupied by anthropogenic activities that may release chemical contaminants (S5 Fig). DWI_1 and DWI_2 were moderately vulnerable to chemical contamination (55.2% and 65% probability in the medium state, respectively). They were also influenced by the land use upstream of the intakes, as illustrated in S5 Fig.

3.3 Probability of the hazard

The findings related to the hazard nodes, which represent another risk component, are presented in this section (Eq 1). The protection area and drainage basin delineation results showed that the study site includes 25 DBs that discharge within the boundaries of the




Fig 5. Bar plots for BN sub-model assessing the probability distribution of overall hazard, crop activity hazard, and animal production hazard within the drainage basins.

<https://doi.org/10.1371/journal.pwat.0000073.g005>

intermediate protection zones of the four DWIs. The immediate protection areas of all intakes and seven drainage basins (DB_1, DB_3, DB_5, DB_16, DB_17, DB_21, and DB_22) were not occupied by agricultural activities, so they do not constitute a risk from agriculture to water quality at DWIs. The remaining drainage basins and their occupancy by animal and crop production activities are shown in the S6–S14 Figs. The minimum areas of crop and animal production activities were 0.08 km² (in DB_20) and 0.03 km² (in DB_12 and DB_18), respectively, while the DB_8 has the greatest area covered by these activities. The crop activities were spread over 3488.26 km², and the animal activities were occupied by 3177.22 km² of this DB. The total area occupying all the DBs upstream of all the DWIs were 8278 km² in crop production activities and 7830 km² in animal activities.

Fig 5 shows the probability distributions of the overall hazard (PH), the crop activity hazard (CAH), and the animal production hazard (APH) for each DB. The probability of hazard was at high level in DB_8, DB_10 and DB_23 (81.7%, 96.6% and 81.6% probability in the high state, respectively). This outcome is due to the area occupied by agricultural activities and the type of production that presents the greatest hazard to water quality: class I for animal production and annual crop for vegetable production. The probability of hazard was moderate in DB_6, where 51.1% was in the medium state. For DB_2, DB_7, DB_15, and DB_24, the probability of hazard was ranged between low and medium levels with close percentages (for example, 48.7% and 49.6% probability in low and medium states, respectively for DB_2). The remaining DBs were all characterised by low hazard probabilities level, which averaged 63.6%.

3.4 Global risk

This section reports the global risk results, which drive from the combination of hazard, vulnerability, and exposure. The overall risk from crop and animal activities was relatively high in three drainage basins (DB_8, DB_10, DB_23) whose outlets were located upstream of these DWIs: DWI_2, DWI_3, and DWI_4 (Fig 6). Their probabilities of being in high status vary between 81.9% and 95.5%. Another five DBs (DB_2, DB_6, DB_7, DB_15, and DB_24) represented a moderate overall risk with probabilities ranging from 55.6% to 60.9% (Fig 6). Eight DBs (DB_4, DB_11, DB_12, DB_13, DB_14, DB_18, DB_19, and DB_25) were associated with




Fig 6. Global risk distribution for all drainage basins upstream four DWIs.

<https://doi.org/10.1371/journal.pwat.0000073.g006>

a low global risk with probabilities that are almost equal between the low and medium status, and with a variance in probabilities that does not exceed 10%. DB_9 and DB_20 were associated with low global risks with probabilities ranging from 54.8 to 67.4% in the low status.

The crops activities hazard (CAH), animal production hazard (APH), exposure (EXP), DWI vulnerability level (DWIVL), vulnerability to microbiological contamination (VMC), and vulnerability to chemical contamination (VCC) are the nodes of interest in this study, so they were selected as findings nodes. Subsequently, the sensitivity for the endpoint "Global Risk Level (GRL)" in every DB occupied by agricultural activities was calculated and analyzed as explained in section 2.3.3. Results of this analysis (Fig 7) showed that the developed model was most sensitive to the crops activities hazard (CAH) (from 17.8 to 29.5% variance




Fig 7. Results of sensitivity of the « Global risk level » variable.

<https://doi.org/10.1371/journal.pwat.0000073.g007>

reduction) and the animal production hazard (APH) (from 6.54 to 25.6%) for all DWIs. This model was also moderately influenced by the exposure node (from 2.5 to 4.13%), especially for the cases of the drainage basins that were slightly occupied by a single type of production activity, either crop or animal, such as BD_9 and BD_20 that are located upstream DWI_2, DWI_3 and DWI_4.

4. Discussion

The probability of DWIs contamination by agricultural activities was assessed using a probabilistic approach in this study. The proposed Bayesian model generates a probabilistic estimation of the DWIs vulnerability level, crop and animal production activities hazard, the exposure, and the overall risk associated with each drainage basin occupied by agricultural activities.

One of the strengths of this model is the ability to embed different types of data in the same system. For example, to fill the input variables, water quality and land use data were used, and the outputs of other studies (vulnerability to water scarcity) can be included. The discretization of all nodes constituting this model allows for more efficient and effective communication with stakeholders on the components of the risk assessment, the interpretation of risks, and the preparation of action plans, since the terminology and statuses of nodes are based on regulatory criteria and lexicons already used in the provincial databases. The graphical model can be used as a communication tool between scientists, municipal officials, government professionals, and members of a watershed organization. It could also be a simple approach for farmer organizations that support surface water pollution reduction to convey potential risks to their members. The acyclic graph facilitates understanding and collaboration among multidisciplinary members, thus increasing the impact in the decision-making process, as discussed in Stritih et al. [98] and Laurila-Pant et al. [99].

In testing several methods to fill CPTs, it was concluded that the chosen methodology can significantly influence the final output of the risk assessment. The additive aggregation was applied through the mixed aggregation by choosing $\alpha = 1$ to fill CPTs related to child nodes of crops activities hazard (CAH), animal production hazard (APH), probability of hazard (PH), and global risk level (GRL). This method is suitable for cases that do not require emphasizing the pessimism bias. For instance, a high level of crop activity (CAL node) may not represent a high hazard (CAH node) in the case where most of the area is covered (CPC node) by undefined crops or non-cultivated land. In another case, the worst-case bias was emphasized by setting α to 0, which applied to the case of assessing vulnerability to chemical contaminants based on land use (CVLU node) in immediate (CRIM node) and intermediate protection zone (CRIN node). This means that the maximum aggregation is applied to penalize the worst of inputs to protect the water resource against the highest hazard to water quality. However, for the rest of the cases, combining both types of aggregation (additive and maximal) was done due to the evaluation of complementary variables, such as the case of chemical and microbiological water quality assessment based on raw water quality analysis. The mixed aggregation method enables the inclusion of inter-correlation between parent nodes, resulting in a risk assessment that is more unbiased and objective compared to other methods. Our case study also revealed that the weight assigned to each parent node might reflect its relative significance and reliance on its respective child node.

As shown in Fig 6 and maps of agricultural activities (from S6–S14 Figs), medium level risk probability dominates for DBs that were slightly occupied by agricultural activities. This was due to the high DWIs vulnerability, and the location of the outlets of the DBs at a distance that does not exceed 10 km upstream from these intakes. This implies that even though the hazard probability is not so significant, the vulnerability of the intake and the exposure of the threat

could be the factors contributing to a higher risk level (medium to high). Therefore, all nodes in the model are essential for assessing the risk of agricultural activities. As a result, across a basic network graph, this Bayesian model gives a clearer understanding of the relative relationship and interaction between all variables [100, 101]. As anticipated, results proved that the BN could be applied to water resource risk assessment with the same efficiency as environmental risk assessment [38]. The developed model provides a transparent and systematic way to characterize risk through uncertainty. It reveals all risk states with their estimated probabilities.

The sensitivity analysis results showed that, in order to improve the global risk estimation, it is most impactful to focus more on the CAH, APH, and EXP since they are linked to the highest entropy reduction. For this purpose, the use of data regarding pesticides (types, quantities, method of application and frequency), animal production, and characteristics of the receiving environment (soil type, hydrodynamics of the river, etc.) are beneficial to have a more accurate assessment. These data are not available in provincial databases. To bridge this data gap, targeted analysis are thus needed. Such models are suitable for determining the catchment areas that could potentially result in the highest risk to water intakes. This simple model is handy for authorities and stakeholders to estimate the level of risk on a regional and provincial scale with commonly available data. Therefore, this model is considered a jump-start for stakeholders to channel their efforts and investments properly to the most appropriate target for contamination mitigation efforts. This risk assessment model may benefit from an upgrade over time by including variables that define the modeling of microbial pathogens and chemical contaminants after rain discharge. As shown by Haack et al. [79], Soupir et al. [76], and Jaffrezic et al. [77], water bodies were most sensitive to livestock-derived fecal and chemical marker inputs after rainfall, and bacteria concentrations frequently exceed standards for primary contact.

Discussion of the risk analysis outcomes with stakeholders requires recognizing uncertainties resulting from the lack of data, inaccuracies in laboratory measurements for raw water quality monitoring, gaps in the knowledge, and inexactness of geographic data. Nevertheless, as Uusitalo [39] pointed out, the advantage of the BN is that it incorporates uncertainty explicitly and naturally through the probability distribution.

As expected, this model has a limitation concerning its validation since it is not readily possible to perform a quantitative validation. This limitation is in accordance with the findings discussed in reviews by Kaikkonen et al. [38] and Phan et al. [51], which also exhibited the same issue. For this reason, in this study, a qualitative model validation was performed by examining the outcomes of many scenarios in comparison to the regulatory vulnerability analyses. Results were consistent; however, the BN approach provides more context for risk levels and an approach to provide an aggregated and global assessment of DWI vulnerability.

This model is a potential tool for source water managers to easily apply since it is based on readily available input data and raw water analyses commonly required by drinking water quality regulations [80], which specify standards for all water systems designated for human use throughout the province of Quebec. Consequently, it is simple to apply this Bayesian model throughout Quebec by modifying the input values in the model to generate probabilistic results regarding the risk posed by agricultural activities upstream of DWIs. In this case, all model components, including parent and child nodes and their probabilistic connections, are not subject to modification. This Bayesian model can also be applied to other study sites on a larger scale (national and international scale) if all the data defining the input nodes are available. As mentioned by Dorner et al. [50], the Bayesian network has the advantage of being used rapidly to investigate outputs under various scenarios. This evidence holds when applying the same Bayesian model that has already been developed. It is crucial to highlight that the groundwork for developing this model, including the generation of CPTs, took a substantial

amount of time and effort. If the necessary data is unavailable for other study sites, adjustments must be made to the model design based on the existing data. As part of these adjustments, it is necessary to decide on the appropriate nodes to use as input to the model while considering the targeted region's regulations. This decision should be made based on the data available, especially those that describe the agricultural activities inventoried at the research site and the quality of surface water at the DWI. The adjustments would also include defining the conditional probability distributions that characterize the probabilistic links of the new nodes added with their child nodes, despite applying the identical CPTs filling approaches described in section 2.2.2. At this scale, the involvement of experts is a must for the qualitative validation of the model.

5. Conclusions

This study used the Bayesian network (BN) to assess the risk of agricultural activities for four DWIs that draw their water across a river in the south of Quebec, Canada. The findings have led us to conclude that:

- The Bayesian model allowed for the combination and manipulation of data in various formats (numerical, statistical, and geographical data) into a consistent methodology for water resource risk assessment for drinking water supplies.
- The interactions between all risk components (the vulnerability of drinking water intake, the exposure to the threat, and the probability of hazard) and their combined uncertainties can be quantified in a single model. Stakeholders and agricultural organizations may utilize this probabilistic reasoning tool to grasp the causal relationships between variables, allowing them to identify the key factors that demand special attention due to their large impacts on the outcomes.
- The end-user can fill data gaps in the model by referencing expert knowledge. The explicit identification of data gaps and sensitivity analyses facilitates the prioritization of future data collection needs. Furthermore, the model can serve as a basis for integrating local knowledge from various stakeholders in the watershed.
- The mixed aggregation method ($f_{add-max}$) combined with expert judgment is a simple and straightforward way for populating conditional probability tables, and it is suitable for considering parent-child dependencies. The correlation between the parent nodes is also considered, making the evaluation accurate based on expert judgment. The weights allocated to the additive (f_{add}) and maximum (f_{max}) aggregation allow for a more focused examination of the influence factor in the field of water resource protection. This is a crucial characteristic when the effect of parent nodes functioning concurrently is considered. This method is useful for filling CPTs semi-automatically to help experts calculate reasonable probabilities when data are not available.
- The model developed in this paper is the first element of a risk assessment cycle. This Bayesian model proved successful in identifying drainage basins occupied by agricultural activities that may pose a medium to high risk of contamination of drinking water intakes. These findings can assist managers and authorities in developing targeted sampling programs for microbiological and chemical testing of surface water while reducing costs and prioritizing drainage basins associated with a high level of risk. The analysis results should constitute the input variables of this Bayesian network while incorporating new nodes and updating it to better pinpoint the source of risk.

- The availability of the data describing the input nodes determines how easily the model may be exported to different water resource risk assessment scenarios. If the data are readily available and suitable for the Bayesian model, the established CPTs in this study can be applied without adjustments. Nevertheless, developing new CPTs would require a lot of time and effort if a new Bayesian network had to be constructed to accommodate other types of input nodes. It is at least plausible to use the same approach of generating conditional probabilities outlined in this study, as it is relevant to the field of water resource protection.

From a water resource protection perspective, in an urban and agricultural watershed, this research is expected to serve as a basis for future studies on assessing microbial risk from combined sewer overflows (CSOs) using a probabilistic approach.

Supporting information

S1 Text. Protection zone delineation.

(DOCX)

S1 Table. Conditional probability distribution.

(DOCX)

S1 Fig. Example of drainage basin (DB) which have outlet located within the boundaries of intermediate protection zone of a drinking water intake.

(DOCX)

S2 Fig. 3D graph of global risk response under the combination of the probability of hazard, exposure, and drinking water intake (DWI) vulnerability level.

(DOCX)

S3 Fig. Detailed Bayesian network structure for assessing drinking water intake contamination risk from agricultural activities.

(DOCX)

S4 Fig. Results of BN submodel assessing the microbiological quality of raw water for all drinking water intakes (DWIs) overall seasons of the year.

(DOCX)

S5 Fig. Results of BN submodel assessing raw water quality for total phosphorus and the chemical vulnerability level based on land use for all drinking water intakes (DWIs).

(DOCX)

S6 Fig. Maps of agricultural activities within the drainage basins (DB_2 and DB_4), as well as the probability distribution of the overall hazard, the crop activity hazard, and the animal production hazard.

(DOCX)

S7 Fig. Maps of agricultural activities within the drainage basins (DB_6 and DB_7), as well as the probability distribution of the overall hazard, the crop activity hazard, and the animal production hazard.

(DOCX)

S8 Fig. Maps of agricultural activities within the drainage basins (DB_8 and DB_9), as well as the probability distribution of the overall hazard, the crop activity hazard, and the animal production hazard.

(DOCX)

S9 Fig. Maps of agricultural activities within the drainage basins (DB_10 and DB_11), as well as the probability distribution of the overall hazard, the crop activity hazard, and the animal production hazard.

(DOCX)

S10 Fig. Maps of agricultural activities within the drainage basins (DB_12 and DB_13), as well as the probability distribution of the overall hazard, the crop activity hazard, and the animal production hazard.

(DOCX)

S11 Fig. Maps of agricultural activities within the drainage basins (DB_14 and DB_15), as well as the probability distribution of the overall hazard, the crop activity hazard, and the animal production hazard.

(DOCX)

S12 Fig. Maps of agricultural activities within the drainage basins (DB_18 and DB_19), as well as the probability distribution of the overall hazard, the crop activity hazard, and the animal production hazard.

(DOCX)

S13 Fig. Maps of agricultural activities within the drainage basins (DB_20 and DB_23), as well as the probability distribution of the overall hazard, the crop activity hazard, and the animal production hazard.

(DOCX)

S14 Fig. Maps of agricultural activities within the drainage basins (DB_24 and DB_25), as well as the probability distribution of the overall hazard, the crop activity hazard, and the animal production hazard.

(DOCX)

Acknowledgments

We thank the partner municipalities and the watershed organization team responsible for the study area for contributing to the data collection and vulnerability assessments.

Author Contributions

Conceptualization: Raja Kammoun, Françoise Bichai.

Data curation: Raja Kammoun.

Formal analysis: Raja Kammoun.

Funding acquisition: Michèle Prévost, Sarah Dorner.

Investigation: Raja Kammoun, Sarah Dorner.

Methodology: Raja Kammoun, Sarah Dorner.

Project administration: Michèle Prévost, Françoise Bichai, Sarah Dorner.

Resources: Michèle Prévost, Françoise Bichai, Sarah Dorner.

Software: Raja Kammoun.

Supervision: Françoise Bichai, Sarah Dorner.

Validation: Raja Kammoun, Natasha McQuaid, Michèle Prévost, Françoise Bichai, Sarah Dorner.

Visualization: Raja Kammoun, Natasha McQuaid.

Writing – original draft: Raja Kammoun.

Writing – review & editing: Raja Kammoun, Natasha McQuaid, Vincent Lessard, Eyerusalem Adhanom Goitom, Michèle Prévost, Françoise Bichai, Sarah Dorner.

References

- Berens MJ, Capel PD, Arnold WA. Neonicotinoid Insecticides in Surface Water, Groundwater, and Wastewater Across Land-Use Gradients and Potential Effects. *Environ Toxicol Chem*. 2021; 40(4):1017–33. <https://doi.org/10.1002/etc.4959> PMID: 33301182
- Deng X, Wagner S, Wang D, Luo Y, Goh KS. Pesticide Detections, Benchmark Exceedances, and Temporal Trends in Surface Water of California's Imperial, Salinas, and Santa Maria Valleys. *Pesticides in Surface Water: Monitoring, Modeling, Risk Assessment, and Management*. ACS Symposium Series. 1308: American Chemical Society; 2019. p. 119–42.
- Geng Y, Jiang L, Zhang D, Liu B, Zhang J, Cheng H, et al. Glyphosate, aminomethylphosphonic acid, and glufosinate ammonium in agricultural groundwater and surface water in China from 2017 to 2018: Occurrence, main drivers, and environmental risk assessment. *Sci Total Environ*. 2021; 769:144396. <https://doi.org/10.1016/j.scitotenv.2020.144396> PMID: 33486182
- Ma L, Yates SR, Ashworth D. Parent and conjugated estrogens and progestagens in surface water of the Santa Ana River: Determination, occurrence, and risk assessment. *Environ Toxicol Chem*. 2016; 35(11):2657–64. <https://doi.org/10.1002/etc.3447> PMID: 27061433
- Malaj E, von der Ohe PC, Grote M, Kuhne R, Mondy CP, Usseglio-Polatera P, et al. Organic chemicals jeopardize the health of freshwater ecosystems on the continental scale. *Proc Natl Acad Sci U S A*. 2014; 111(26):9549–54. <https://doi.org/10.1073/pnas.1321082111> PMID: 24979762
- Syafrudin M, Kristanti RA, Yuniarto A, Hadibarata T, Rhee J, Al-Onazi WA, et al. Pesticides in Drinking Water-A Review. *Int J Environ Res Public Health*. 2021; 18(2). <https://doi.org/10.3390/ijerph18020468> PMID: 33430077
- Schindler DW, Dillon PJ, Schreier H. A review of anthropogenic sources of nitrogen and their effects on Canadian aquatic ecosystems. In: Martinelli LA, Howarth RW, editors. *Nitrogen Cycling in the Americas: Natural and Anthropogenic Influences and Controls*. Dordrecht: Springer, Dordrecht; 2006. p. 25–44.
- Silva-Madera RJ, Salazar-Flores J, Peregrina-Lucano AA, Mendoza-Michel J, Ceja-Galvez HR, Rojas-Bravo D, et al. Pesticide Contamination in Drinking and Surface Water in the Cienega, Jalisco, Mexico. *Water Air Soil Poll*. 2021; 232(2).
- Ware GW, Albert LA, Voogt P, Gerba CP, Hutzinger O, Knaak JB, et al. *Reviews of Environmental Contamination and Toxicology*. NY, USA: Springer New York, NY; 2006. 261 p.
- Triegel EK, Guo L. Overview of the Fate of Pesticides in the Environment, Water Balance; Runoff vs. Leaching. *Mechanisms Of Pesticide Movement Into Ground Water*. Mechanisms of Pesticide Movement into Ground Water: CRC Press; 2018. p. 1–13.
- Andrade VS, Gutierrez MF, Regaldo L, Paira AR, Repetti MR, Gagneten AM. Influence of rainfall and seasonal crop practices on nutrient and pesticide runoff from soybean dominated agricultural areas in Pampean streams, Argentina. *Sci Total Environ*. 2021; 788:147676. <https://doi.org/10.1016/j.scitotenv.2021.147676> PMID: 34029815
- Harrison S, McAree C, Mulville W, Sullivan T. The problem of agricultural 'diffuse' pollution: Getting to the point. *Sci Total Environ*. 2019; 677:700–17. <https://doi.org/10.1016/j.scitotenv.2019.04.169> PMID: 31071672
- Kuo JN, Soon AY, Garrett C, Wan MT, Pasternak JP. Agricultural pesticide residues of farm runoff in the Okanagan Valley, British Columbia, Canada. *J Environ Sci Health B*. 2012; 47(4):250–61. <https://doi.org/10.1080/03601234.2012.636588> PMID: 22428886
- Calliera M, Capri E, Zambito Marsala R, Russo E, Bisagni M, Colla R, et al. Multi-actor approach and engagement strategy to promote the adoption of best management practices and a sustainable use of pesticides for groundwater quality improvement in hilly vineyards. *Sci Total Environ*. 2021; 752:142251. <https://doi.org/10.1016/j.scitotenv.2020.142251> PMID: 33207509

15. Cahn Michael D, Phillips B. Best Management Practices for Mitigating Pesticides in Runoff from Vegetable Systems in California. *Pesticides in Surface Water: Monitoring, Modeling, Risk Assessment, and Management*. ACS Symposium Series. 1308: American Chemical Society; 2019. p. 519–39.
16. Fisher IJ, Phillips PJ, Bayraktar BN, Chen S, McCarthy BA, Sandstrom MW. Pesticides and their degradates in groundwater reflect past use and current management strategies, Long Island, New York, USA. *Sci Total Environ*. 2021; 752:141895. <https://doi.org/10.1016/j.scitotenv.2020.141895> PMID: 32892047
17. Bishop SL, Kerkovius JK, Menard F, Murch SJ. N-beta-Methylamino-L-Alanine and Its Naturally Occurring Isomers in Cyanobacterial Blooms in Lake Winnipeg. *Neurotox Res*. 2018; 33(1):133–42.
18. El-Shehawy R, Gorokhova E, Fernandez-Pinas F, del Campo FF. Global warming and hepatotoxin production by cyanobacteria: what can we learn from experiments? *Water Res*. 2012; 46(5):1420–9. <https://doi.org/10.1016/j.watres.2011.11.021> PMID: 22178305
19. Jankowiak J, Hattenrath-Lehmann T, Kramer BJ, Ladds M, Gobler CJ. Deciphering the effects of nitrogen, phosphorus, and temperature on cyanobacterial bloom intensification, diversity, and toxicity in western Lake Erie. *Limnology and Oceanography*. 2019; 64(3):1347–70.
20. Yeager N, Carpenter A. State approaches to addressing cyanotoxins in drinking water. *AWWA Water Science*. 2019; 1(1).
21. Jalili F, Trigui H, Guerra Maldonado JF, Dorner S, Zamyadi A, Shapiro BJ, et al. Can Cyanobacterial Diversity in the Source Predict the Diversity in Sludge and the Risk of Toxin Release in a Drinking Water Treatment Plant? *Toxins (Basel)*. 2021; 13(1).
22. Shang L, Feng M, Xu X, Liu F, Ke F, Li W. Co-Occurrence of Microcystins and Taste-and-Odor Compounds in Drinking Water Source and Their Removal in a Full-Scale Drinking Water Treatment Plant. *Toxins (Basel)*. 2018; 10(1).
23. Zamyadi A, Dorner S, Sauve S, Ellis D, Bolduc A, Bastien C, et al. Species-dependence of cyanobacteria removal efficiency by different drinking water treatment processes. *Water Res*. 2013; 47(8):2689–700. <https://doi.org/10.1016/j.watres.2013.02.040> PMID: 23515107
24. Zamyadi A, MacLeod SL, Fan Y, McQuaid N, Dorner S, Sauve S, et al. Toxic cyanobacterial breakthrough and accumulation in a drinking water plant: a monitoring and treatment challenge. *Water Res*. 2012; 46(5):1511–23. <https://doi.org/10.1016/j.watres.2011.11.012> PMID: 22137293
25. Maghsoudi E, Fortin N, Greer C, Duy SV, Fayad P, Sauve S, et al. Biodegradation of multiple microcystins and cylindrospermopsin in clarifier sludge and a drinking water source: Effects of particulate attached bacteria and phycocyanin. *Ecotoxicol Environ Saf*. 2015; 120:409–17. <https://doi.org/10.1016/j.ecoenv.2015.06.001> PMID: 26122734
26. Moradinejad S, Trigui H, Guerra Maldonado JF, Shapiro J, Terrat Y, Zamyadi A, et al. Diversity Assessment of Toxic Cyanobacterial Blooms during Oxidation. *Toxins (Basel)*. 2020; 12(11). <https://doi.org/10.3390/toxins12110728> PMID: 33233813
27. Dufour AP, Bartram JK, Bos R, Gannon VPJ. *Animal Waste, Water Quality and Human Health*. London, UK: World Health Organisation (WHO) WHO & United States Environmental Protection Agency (USEPA); 2012.
28. Wilkes G, Edge T, Gannon V, Jokinen C, Lyautey E, Medeiros D, et al. Seasonal relationships among indicator bacteria, pathogenic bacteria, *Cryptosporidium* oocysts, *Giardia* cysts, and hydrological indices for surface waters within an agricultural landscape. *Water Res*. 2009; 43(8):2209–23. <https://doi.org/10.1016/j.watres.2009.01.033> PMID: 19339033
29. Sylvestre E, Burnet JB, Smeets P, Medema G, Prevost M, Dorner S. Can routine monitoring of *E. coli* fully account for peak event concentrations at drinking water intakes in agricultural and urban rivers? *Water Res*. 2020; 170:115369.
30. Edge TA, El-Shaarawi A, Gannon V, Jokinen C, Kent R, Khan IU, et al. Investigation of an *Escherichia coli* environmental benchmark for waterborne pathogens in agricultural watersheds in Canada. *J Environ Qual*. 2012; 41(1):21–30. <https://doi.org/10.2134/jeq2010.0253> PMID: 22218170
31. Alegbeleye OO, Sant'Ana AS. Manure-borne pathogens as an important source of water contamination: An update on the dynamics of pathogen survival/transport as well as practical risk mitigation strategies. *Int J Hyg Environ Health*. 2020; 227:113524. <https://doi.org/10.1016/j.ijheh.2020.113524> PMID: 32298989
32. Bonansea RI, Filippi I, Wunderlin DA, Marino DJG, Ame MV. The Fate of Glyphosate and AMPA in a Freshwater Endorheic Basin: An Ecotoxicological Risk Assessment. *Toxics*. 2017; 6(1).
33. Carazo-Rojas E, Perez-Rojas G, Perez-Villanueva M, Chinchilla-Soto C, Chin-Pampillo JS, Aguilar-Mora P, et al. Pesticide monitoring and ecotoxicological risk assessment in surface water bodies and sediments of a tropical agro-ecosystem. *Environ Pollut*. 2018; 241:800–9. <https://doi.org/10.1016/j.envpol.2018.06.020> PMID: 29909306

34. Holmes CM, Brown CD, Hamer M, Jones R, Maltby L, Posthuma L, et al. Prospective aquatic risk assessment for chemical mixtures in agricultural landscapes. *Environ Toxicol Chem*. 2018; 37(3):674–89. <https://doi.org/10.1002/etc.4049> PMID: 29193235
35. Jiang YF, Chen SC, Hu BF, Zhou Y, Liang ZZ, Jia XL, et al. A comprehensive framework for assessing the impact of potential agricultural pollution on grain security and human health in economically developed areas. *Environmental Pollution*. 2020; 263.
36. Tan H, Zhang H, Wu C, Wang C, Li Q. Pesticides in surface waters of tropical river basins draining areas with rice-vegetable rotations in Hainan, China: Occurrence, relation to environmental factors, and risk assessment. *Environ Pollut*. 2021; 283:117100. <https://doi.org/10.1016/j.envpol.2021.117100> PMID: 33865099
37. Tokatli C. Health risk assessment of toxic metals in surface and groundwater resources of a significant agriculture and industry zone in Turkey. *Environ Earth Sci*. 2021; 80(4).
38. Kaikkonen L, Parviainen T, Rahikainen M, Uusitalo L, Lehtikainen A. Bayesian Networks in Environmental Risk Assessment: A Review. *Integr Environ Assess Manag*. 2020; 17(1):62–78. <https://doi.org/10.1002/ieam.4332> PMID: 32841493
39. Uusitalo L. Advantages and challenges of Bayesian networks in environmental modelling. *Ecological Modelling*. 2007; 203(3–4):312–8.
40. Harris MJ, Stinson J, Landis WG. A Bayesian Approach to Integrated Ecological and Human Health Risk Assessment for the South River, Virginia Mercury-Contaminated Site. *Risk Anal*. 2017; 37(7):1341–57. <https://doi.org/10.1111/risa.12691> PMID: 28121045
41. McDonald KS, Ryder DS, Tighe M. Developing best-practice Bayesian Belief Networks in ecological risk assessments for freshwater and estuarine ecosystems: a quantitative review. *J Environ Manage*. 2015; 154:190–200. <https://doi.org/10.1016/j.jenvman.2015.02.031> PMID: 25733196
42. O'Brien GC, Dickens C, Hines E, Wepener V, Stassen R, Quayle L, et al. A regional-scale ecological risk framework for environmental flow evaluations. *Hydrology and Earth System Sciences*. 2018; 22(2):957–75.
43. Shan K, Shang M, Zhou B, Li L, Wang X, Yang H, et al. Application of Bayesian network including *Microcystis* morphospecies for microcystin risk assessment in three cyanobacterial bloom-plagued lakes, China. *Harmful Algae*. 2019; 83:14–24. <https://doi.org/10.1016/j.hal.2019.01.005> PMID: 31097252
44. Liu KF-R, Lu C-F, Chen C-W, Shen Y-S. Applying Bayesian belief networks to health risk assessment. *Stochastic Environmental Research and Risk Assessment*. 2011; 26(3):451–65.
45. Sperotto A, Molina JL, Torresan S, Critto A, Marcomini A. Reviewing Bayesian Networks potentials for climate change impacts assessment and management: A multi-risk perspective. *J Environ Manage*. 2017; 202(Pt 1):320–31. <https://doi.org/10.1016/j.jenvman.2017.07.044> PMID: 28750284
46. Li L, Wang J, Leung H, Jiang C. Assessment of catastrophic risk using Bayesian network constructed from domain knowledge and spatial data. *Risk Anal*. 2010; 30(7):1157–75. <https://doi.org/10.1111/j.1539-6924.2010.01429.x> PMID: 20497388
47. Li M, Hong M, Zhang R. Improved Bayesian Network-Based Risk Model and Its Application in Disaster Risk Assessment. *International Journal of Disaster Risk Science*. 2018; 9(2):237–48.
48. Zhu ZJY, Xiang Y, McBean E. Operational Hazard Risk Assessment Using Bayesian Networks. *Iceis 2011: Proceedings of the 13th International Conference on Enterprise Information Systems, Vol 2*. 2011:135–9.
49. Yu R, Zhang C. Early warning of water quality degradation: A copula-based Bayesian network model for highly efficient water quality risk assessment. *J Environ Manage*. 2021; 292:112749. <https://doi.org/10.1016/j.jenvman.2021.112749> PMID: 34004503
50. Dorner S, Shi J, Swayne D. Multi-objective modelling and decision support using a Bayesian network approximation to a non-point source pollution model. *Environmental Modelling & Software*. 2007; 22(2):211–22.
51. Phan TD, Smart JCR, Capon SJ, Hadwen WL, Sahin O. Applications of Bayesian belief networks in water resource management: A systematic review. *Environmental Modelling & Software*. 2016; 85:98–111.
52. Jalliffier-Verne I, Leconte R, Huaranga-Alvarez U, Madoux-Humery AS, Galarneau M, Servais P, et al. Impacts of global change on the concentrations and dilution of combined sewer overflows in a drinking water source. *Sci Total Environ*. 2015; 508:462–76. <https://doi.org/10.1016/j.scitotenv.2014.11.059> PMID: 25506909
53. Werstuck C, Coulibaly P. Hydrometric network design using dual entropy multi-objective optimization in the Ottawa River Basin. *Hydrology Research*. 2017; 48(6):1639–51.

54. Government of Quebec. Water Withdrawal and Protection Regulation- Q-2, r. 35.2. Updated to June 1, 2022 ed. Quebec, Canada: Quebec Official Publisher; 2014. p. 48.
55. Pearl J. Fusion, Propagation, and Structuring in Belief Networks. *Artificial Intelligence*. 1986; 29(3):241–88.
56. Bozorgi A, Roozbahani A, Shahdany SMH, Abbassi R. Development of Multi-Hazard Risk Assessment Model for Agricultural Water Supply and Distribution Systems Using Bayesian Network. *Water Resources Management*. 2021; 35(10):3139–59.
57. Barton DN, Sundt H, Bustos AA, Fjeldstad HP, Hedger R, Forseth T, et al. Multi-criteria decision analysis in Bayesian networks—Diagnosing ecosystem service trade-offs in a hydropower regulated river. *Environmental Modelling & Software*. 2020; 124.
58. Bertone E, Sahin O, Richards R, Roiko A. Extreme events, water quality and health: A participatory Bayesian risk assessment tool for managers of reservoirs. *Journal of Cleaner Production*. 2016; 135:657–67.
59. Keshtkar AR, Salajegheh A, Sadoddin A, Allan MG. Application of Bayesian networks for sustainability assessment in catchment modeling and management (Case study: The Hablehrood river catchment). *Ecological Modelling*. 2013; 268:48–54.
60. Grêt-Regamey A, Straub D. Spatially explicit avalanche risk assessment linking Bayesian networks to a GIS. *Natural Hazards and Earth System Sciences*. 2006; 6(6):911–26.
61. Norsys Software and Corp. Netica, Version 6.09 2021 [10 november 2021]. Available from: <https://www.norsys.com/netica.html>.
62. Chen Y-C, Wheeler TA, Kochenderfer MJ. Learning Discrete Bayesian Networks from Continuous Data. *Journal of Artificial Intelligence Research*. 2017; 59:103–32.
63. Intergovernmental Panel on Climate Change. Climate Change 2022: Impacts, Adaptation and Vulnerability. Summary for Policymakers. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge, UK and New York, NY, USA: Cambridge University Press; 2022. p. 3676.
64. World Health Organization (WHO). Protecting Surface Water for Health. Identifying, Assessing and Managing Drinking-Water Quality Risks in Surface-Water Catchments. Geneva; 2016. Report No.: ISBN 978 92 4 151055 4.
65. Kammoun R, Leveque B, Prévost M, Bichai F, Dorner S. Indice de vulnérabilité d'un site de prélèvement d'eau de surface: étude de cas d'un bassin versant urbain et agricole situé dans le sud du Québec. *Révue Génial*. 2020.
66. Government of Canada. When do the seasons start? Canada: Government of Canada; 2021 [15 December 2021]. Available from: <https://nrc.canada.ca/en/certifications-evaluations-standards/canadas-official-time/3-when-do-seasons-start>.
67. Leveque B. Analyse des vulnérabilités des prises d'eau potable de la rivière des Mille-Îles (Québec) aux étiages estivaux en contexte de changements globaux par une approche ascendante. Montréal Polytechnique Montréal; 2020.
68. Ministère de l'Environnement et de la Lutte contre les Changements climatiques (MELCC). Portail des connaissances sur l'eau (PCE) Canada2022 [15 December 2021]. Available from: <http://www.environnement.gouv.qc.ca/eau/portail/>.
69. Government of Quebec. Land Use Mapping of the St. Lawrence Lowlands Canada2019 [15 December 2021]. Available from: <https://www.donneesquebec.ca/recherche/fr/dataset/cartographie-de-l-occupation-du-sol-des-basses-terres-du-saint-laurent>.
70. Beaudin I, Deslandes J, Michaud AR, Bonn F, Madramootoo CA. Variabilité spatio-temporelle des exportations de sédiments et de phosphore dans le bassin versant de la Rivière aux Brochets au sud-ouest du Québec—Partie I: Paramétrage, calibrage et validation du modèle SWAT. *Agrosolutions*. 2006:4–20.
71. Michaud A, Deslandes J, Beaudin I. Modélisation de l'hydrologie et des dynamiques de pollution diffuse dans le bassin-versant de la rivière aux Brochets à l'aide du modèle SWAT. Canada: Institut de Recherche et de Développement en Agroenvironnement (IRDA); 2006.
72. Gasser MO, Bolinder M, Martel S, Poulin D, Beaudin I, Michaud A, et al. Impacts agroenvironnementaux associés à la culture et au prélèvement de biomasses végétales agricoles pour la production de bioproduits industriels. Rapport final. Canada: Institut de recherche et de développement en agroenvironnement; 2011.
73. Giroux I. Présence de pesticides dans l'eau au Québec: Portrait et tendances dans les zones de maïs et de soya—2011 à 2014. Québec, Canada: Ministère du Développement durable, de l'Environnement et de la Lutte contre les changements climatiques, Direction du suivi de l'état de l'environnement; 2015.

74. Giroux I. Présence de pesticides dans l'eau au Québec: Portrait et tendances dans les zones de maïs et de soya— 2015 à 2017. Québec: Ministère de l'Environnement et de la Lutte contre les changements climatiques, Direction générale du suivi de l'état de l'environnement; 2019.
75. Government of Quebec. Land use Canada2017 [15 December 2021]. Available from: <https://www.donneesquebec.ca/recherche/fr/dataset/utilisation-du-territoire>.
76. Soupir ML, Mostaghimi S, Yagow ER, Hagedorn C, Vaughan DH. Transport of fecal bacteria from poultry litter and cattle manures applied to pastureland. *Water Air Soil Poll.* 2006; 169(1–4):125–36.
77. Jaffrezic A, Jarde E, Pourcher AM, Gourmelon M, Caprais MP, Heddadji D, et al. Microbial and chemical markers: runoff transfer in animal manure-amended soils. *J Environ Qual.* 2011; 40(3):959–68. <https://doi.org/10.2134/jeq2010.0355> PMID: 21546682
78. Lalancette C, Genereux M, Mailly J, Servais P, Cote C, Michaud A, et al. Total and infectious Cryptosporidium oocyst and total Giardia cyst concentrations from distinct agricultural and urban contamination sources in Eastern Canada. *J Water Health.* 2012; 10(1):147–60. <https://doi.org/10.2166/wh.2011.049> PMID: 22361710
79. Haack SK, Duris JW, Kolpin DW, Focazio MJ, Meyer MT, Johnson HE, et al. Contamination with bacterial zoonotic pathogen genes in U.S. streams influenced by varying types of animal agriculture. *Sci Total Environ.* 2016; 563–564:340–50. <https://doi.org/10.1016/j.scitotenv.2016.04.087> PMID: 27139306
80. Government of Quebec. Regulation respecting the quality of drinking water-Q-2, r. 40 (Environment Quality Act). Updated to June 1, 2022 ed. Québec: Québec Official Publisher; 2012. p. 68.
81. Hébert S. Développement d'un indice de la qualité bactériologique et physico-chimique de l'eau pour les rivières du Québec. Québec: Ministère de l'Environnement et de la Faune; 1997. Report No.: EN/970102.
82. Kammoun R, Lessard V, Gagnière C, Bichai F, Prévost M, Goulet R, et al. Rapport d'analyse de la vulnérabilité de la source pour le prélèvement d'eau de surface no. X1300278-1. Montréal, QC, Canada: Polytechnique Montréal & COBAMIL; 2021.
83. Kammoun R, Lessard V, Gagnière C, Bichai F, Prévost M, Goulet R, et al. Rapport d'analyse de la vulnérabilité de la source pour le prélèvement d'eau de surface no. 24846859–1. Montréal, QC, Canada: Polytechnique Montréal & COBAMIL 2021.
84. Kammoun R, Lessard V, Gagnière C, Bichai F, Prévost M, Goulet R, et al. Rapport d'analyse de la vulnérabilité de la source pour les prélèvements d'eau de surface no. X0008485-1 et X0008485-2. Montréal, QC, Canada: Polytechnique Montréal & COBAMIL 2021.
85. Kammoun R, Lessard V, Gagnière C, Bichai F, Prévost M, Goulet R, et al. Rapport d'analyse de la vulnérabilité de la source pour les prélèvements d'eau de surface no. X0008104-1 et X0008104-2. Montréal, QC, Canada: Polytechnique Montréal & COBAMIL 2021.
86. Kammoun R, Lessard V, Gagnière C, Bichai F, Prévost M, Goulet R, et al. Rapport d'analyse de la vulnérabilité de la source pour le prélèvement d'eau de surface no. X0008125-1. Montréal, QC, Canada: Polytechnique Montréal & COBAMIL 2021.
87. Kammoun R, Lessard V, Gagnière C, Bichai F, Prévost M, Goulet R, et al. Rapport d'analyse de la vulnérabilité de la source pour le prélèvement d'eau de surface no. X0010573-1. Montréal, QC, Canada: Polytechnique Montréal & COBAMIL 2021.
88. Das B. Generating Conditional Probabilities for Bayesian Networks: Easing the Knowledge Acquisition Problem. *CoRR* 1–24. ArXiv. 2004: 24.
89. Cain J. Planning improvements in natural resource management. Guidelines for using Bayesian networks to support the planning and management of development programmes in the water sector and beyond. Crowmarsh Gifford, Wallingford, Oxon, U K: Centre for Ecology & Hydrology; 2001.
90. Mkrtchyan L, Podofilini L, Dang VN. Methods for building Conditional Probability Tables of Bayesian Belief Networks from limited judgment: An evaluation for Human Reliability Application. *Reliability Engineering & System Safety.* 2016; 151:93–112.
91. Ban SS, Pressey RL, Graham NAJ. Assessing interactions of multiple stressors when data are limited: A Bayesian belief network applied to coral reefs. *Global Environmental Change.* 2014; 27:64–72.
92. Mkrtchyan L, Podofilini L, Dang VN. Overview of methods to build Conditional Probability Tables with partial expert information for Bayesian Belief Networks. In: Group TF, editor. *Safety and Reliability of Complex Engineered Systems.* London2015.
93. Langhans SD, Reichert P, Schuwirth N. The method matters: A guide for indicator aggregation in ecological assessments. *Ecological Indicators.* 2014; 45:494–507.
94. Pearl J. Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference. 340 Pine Street, Sixth Floor, San Francisco, CA, United States: Morgan Kaufmann Publishers Inc.; 1988. 552 p.

95. Pollino CA, Woodberry O, Nicholson A, Korb K, Hart BT. Parameterisation and evaluation of a Bayesian network for use in an ecological risk assessment. *Environmental Modelling & Software*. 2007; 22(8):1140–52.
96. Marcot BG, Steventon JD, Sutherland GD, McCann RK. Guidelines for developing and updating Bayesian belief networks applied to ecological modeling and conservation. *Canadian Journal of Forest Research*. 2006; 36(12):3063–74.
97. Marcot BG. Metrics for evaluating performance and uncertainty of Bayesian network models. *Ecological Modelling*. 2012; 230:50–62.
98. Stritih A, Rabe SE, Robaina O, Gret-Regamey A, Celio E. An online platform for spatial and iterative modelling with Bayesian Networks. *Environmental Modelling & Software*. 2020; 127.
99. Laurila-Pant M, Mantyniemi S, Venesjarvi R, Lehtikoinen A. Incorporating stakeholders' values into environmental decision support: A Bayesian Belief Network approach. *Sci Total Environ*. 2019; 697:134026. <https://doi.org/10.1016/j.scitotenv.2019.134026> PMID: 31476493
100. Kanen R, Marengo MCR, Abdel-Moati H, Cranefield J, Vechot L. Developing a framework for dynamic risk assessment using Bayesian networks and reliability data. *Journal of Loss Prevention in the Process Industries*. 2017; 50:142–53.
101. Mentzel S, Grung M, Tollefsen KE, Stenrød M, Petersen K, Moe SJ. Development of a Bayesian network for probabilistic risk assessment of pesticides. *bioRxiv*. 2021: 2021.05.20.444913.