

Titre: Title:	Implementation of the RESKR module in NJOY
Auteurs: Authors:	Tiejun Zu, Alain Hébert, Liangzhi Cao, & Hongchun Wu
Date:	2022
Type:	Communication de conférence / Conference or Workshop Item
Référence: Citation:	Zu, T., Hébert, A., Cao, L., & Wu, H. (2022, July). Implementation of the RESKR module in NJOY [Paper]. 15th International Conference on Nuclear Data for Science and Technology (ND2022) (2 pages). Published in EPJ Web of Conferences, 284. https://doi.org/10.1051/epjconf/202328408014

 **Document en libre accès dans PolyPublie**

Open Access document in PolyPublie

URL de PolyPublie: PolyPublie URL:	https://publications.polymtl.ca/55702/
Version:	Version officielle de l'éditeur / Published version Révisé par les pairs / Refereed
Conditions d'utilisation: Terms of Use:	CC BY

 **Document publié chez l'éditeur officiel**

Document issued by the official publisher

Nom de la conférence: Conference Name:	15th International Conference on Nuclear Data for Science and Technology (ND2022)
Date et lieu: Date and Location:	2022-07-21 - 2022-07-29
Maison d'édition: Publisher:	EDP Sciences
URL officiel: Official URL:	https://doi.org/10.1051/epjconf/202328408014
Mention légale: Legal notice:	

Implementation of the RESKR module in NJOY

Tiejun Zu^{1,*}, Alain Hébert^{2,**}, and Liangzhi Cao^{1,***} Hongchun Wu^{1,****}

¹School of Nuclear Science and Technology
Xi'an Jiaotong University
China

²Polytechnique Montreal
PO Box 6079, Station Centre-Ville
Montreal QC, Canada, H3C 3A7

Abstract. The Blackshaw-Murray[1] elastic kernel represents the effect of neutron up-scattering caused by thermal motion of target nuclei and resonance elastic scattering on the multigroup scattering matrices. A first implementation of this model was proposed by Ouisloumen and Sanchez and made available in proprietary cross-sections libraries of the PARAGON and APOLLO lattice codes.[2] Later, the same technique was reimplemented as module RESK in the NECP-Atlas cross-section generating code by J. Xu, T. Zu and L. Cao.[3, 4] The proposed RESKR module in NJOY-2012 and NJOY-2016 is based on the implementation of module RESK in the NECP-Atlas cross-section generating code.[5] The RESKR module is released under the BSD Open-Source license.

1 Introduction

The new RESKR module of NJOY generates Doppler-broadened resonance elastic scattering kernels (RESKs) in PENDF format starting from piecewise linear cross sections also in PENDF format. The input OK cross sections are recovered from RECONR and from a previous BROADR run. The RESKR module is dedicated to deterministic calculations and requires further processing using the GROUPR module so as to produce the output multigroup cross-section data, as depicted in Fig. 1.

The exact Doppler broadened energy transfer kernel produces a strong up-scattering effect in the resolved resonance energy range, above the upper limit where the $S(\alpha, \beta)$ free gas model of the THERMR module is set. The RESKR module is used to pursue Doppler broadening of the elastic kernel for resonant nuclides above the thermal domain considered by THERMR.

The RESKR module implements the Blackshaw-Murray kernel according to the Ouisloumen and Sanchez formulas. Firstly, the resonance elastic scattering kernel (RESK) formulations for anisotropic scattering up to any Legendre order is adopted to represent the exact Doppler broadened energy transfer kernels. A semi-analytical integration method is applied to perform the RESK calculations. Combining with the RESK calculations, a linearization algorithm is proposed to generate the RESK interpolation tables. These interpolation tables are Legendre moments ℓ of the elastic scattering kernels of the form $P_\ell(E \rightarrow E')$.

*e-mail: tiejun@xjtu.edu.cn

**e-mail: alain.hebert@polymtl.ca

***e-mail: caolz@mail.xjtu.edu.cn

****e-mail: hongchun@mail.xjtu.edu.cn

They are written in the output PENDF file as a new MF6 MT300 reaction type.

The main entry point is subroutine reskr exported by the Fortran-90 module reskm. The coding logic of module reskm is similar to the one used by the Fortran-90 module thermm.

An unassigned reaction type number MT300 in ENDF-6 format is assigned to store and output the interpolation table of the RESK data and the MF3 MT300 and MF6 MT300 reaction types are defined. The incident energy grid of reaction MF3 MT300 is a subset of energy grid in MF3 MT2 selected between lower (e1o) and upper (ehi) incident energy boundaries for the RESK calculation. The incident energy grid of reaction MF6 MT300 is a coarser grid set to reduce computing cost.

2 The new RESKR module of NJOY

At each reconstructed incident energy, the moments of energy transfer kernels for the different orders are linearized simultaneously on a single unionized grid by the conventional interval-halving techniques of Cullen[6]. It ensures that all orders of the moments are reconstructed smoothly. Meanwhile, the moments of energy transfer kernels for the different orders are interpolated simultaneously once the interpolation interval is found.

A data structure in ENDF-6 format is defined as described in Table 1 where HEAD, TAB2, TAB1 and LIST are the standard types of records; ZA and AWR are the standard material charge and mass parameters; NL is the maximum Legendre order number of this table; T is the absolute temperature; E1 is the primary energy; EP is the




Figure 1. Flow chart of modules in NJOY

Table 1. MF6 MT300 reaction type.

```
[MAT, 6, MT / ZA, AWR, NL, 0, 0, 0 ] HEAD
[MAT, 6, MT / T, 0.0, 0, 0, NR, NE / (INT) ] TAB2
[MAT, 6, MT / 0.0, E1, 0, 0, NRP, NEP / EP / P0(E1->EP) ] TAB1
[MAT, 6, MT / 0.0, 0.0, 0, 0, NW, 0 / P1(E1->EP) ] LIST
Repeat the LIST structure for all NL-1 orders
Repeat the TAB1 structure for all NE incident energies
```

secondary energy; P_0 indicates the 0th moment of energy transfer kernel and P_1 represents the 1st moment of energy transfer kernel. The TAB1 structure with embedded LIST structures is repeated for all NE incident energies. Each TAB1 structure contains NL-1 embedded LIST structures containing kernel values for P_1 , P_2 and higher Legendre moments.

After the processing, the data will be output into derived files which are the point-ENDF (PENDF) files. The reusable PENDF files can be used for generating the different multi-group cross sections and scattering matrices faced with the different requirements of dilutions and energy group structures.

An update of module GROUPR is also required. Subroutine `getrsk` retrieves Legendre-energy data corresponding to the resonance escape scattering kernel and available in the MT300 reaction of the PENDF tape. Subroutine `getrsk` follows the same logic as subroutine `getaed` for thermal scattering data to evaluate the feed function at various incident neutron energies over the panel.

Two multigroup cross section libraries are currently supported: The MATXS format, generated using the MATXSR module and the DRAGLIB format supported by the DRAGR module. The chaining of NJOY modules is depicted in Fig. 1.

3 Conclusion

- The *resonance elastic scattering kernel* (RESK) model is an implementation of Blackshaw-Murray and Sanchez-Ouisloumen elastic kernel consistent with multigroup deterministic calculations.

- The RESK model is the deterministic equivalent of the *Doppler broadening rejection correction* (DBRC) method available in most Monte-Carlo codes.
- Availability of module RESKR is a long-standing request from NJOY community.
- The RESKR module in NJOY is based on the implementation of module RESK in the NECP-Atlas cross-section generating code.
- The new module RESKR is an Open-Source contribution distributed under the BSD license.
- The actual implementation is a beta version requiring further validation. Collaboration is welcome.

References

- [1] G. L. Blackshaw and R. L. Murray, "Scattering functions for low-energy neutron collisions in a Maxwellian monoatomic gas," *Nucl. Sci. Eng.*, **27**, 520–532 (1967).
- [2] M. Ouisloumen and R. Sanchez, "A Model for Neutron Scattering of Heavy Isotopes that Accounts for Thermal Agitation Effects," *Nucl. Sci. Eng.*, **107**, 189–200 (1991).
- [3] T. Zu, J. Xu, Y. Tang, H. Bi, F. Zhao, L. Cao and H. Wu, "NECP-Atlas: A new nuclear data processing code," *Ann nucl. Energy*, **123**, 153–161 (2019).
- [4] J. Xu, T. Zu and L. Cao, "Development and verification of resonance elastic scattering kernel processing module in nuclear data processing code NECP-Atlas," *Progress nucl. Energy*, **110**, 301–310 (2019).
- [5] R. E. MacFarlane and A. C. Kahler, "Methods for Processing ENDF/B-VII with NJOY," *Nuclear Data Sheets*, **111**, 2739 (2010).
- [6] D. E. Cullen, "Program SIGMA1 (Version 74-1)," Lawrence Livermore Laboratory report UCID-16426 (January 1974).